

# ABSOLUTE AQUEOUS REDOX POTENTIALS

(via a new link between aqueous and gaseous properties)

**Raji Heyrovská**

*Institute of Biophysics, Academy of Sciences of the Czech Republic,  
135 Kralovopolská, 612 65 Brno, Czech Rep. Email: rheyrovs@hotmail.com*

Presented at:

**9. Pracovní setkání fyzikálních chemiků a elektrochemiků, Brno, 29. a 30. června 2009** (<http://www.chemi.muni.cz/~libuse/setkani2009/index.html>)

**(9<sup>th</sup> Workshop of Physical Chemists and Electrochemists, Brno, 29th and 30th June 2009)**

on the occasion of the 3 main anniversaries,

- 1) **90<sup>th</sup>**: foundation of Masaryk Univ. (MUNI),
- 2) **90<sup>th</sup>**: foundation of Mendel Univ. of Agriculture & Forestry (MUAF),
- 3) **50<sup>th</sup>**: award of the Nobel Prize in 195**9** to Professor J. Heyrovský,

and also 3 more anniversaries,

- 4) **70<sup>th</sup>**: survival of MUNI & MUAF in 193**9**,
- 5) **40<sup>th</sup>**: survival of MUNI & MUAF again in 196**9** and
- 6) **20<sup>th</sup>**: “ velvet revolution” in 198**9**.

\*\*\*\*\*

# 1. INTRODUCTION

“ **Chemistry is electric**” (Bk: M.E. Bowden, CHF, 1997)

**Tomato and Lemon batteries:**

[http://www.funsci.com/fun3\\_en/electro/electro.htm](http://www.funsci.com/fun3_en/electro/electro.htm) <http://www.youtube.com/watch?v=AY9qcDCFeVI>

**Electrolyte:** acidic juice, **Electrodes:** copper (+) and zinc (-).

**Chemical reactions at the two half cells:** (1)  $\text{Cu}^{++} + 2\text{e}^- \rightleftharpoons \text{Cu}$  and (2)  $\text{Zn} \rightleftharpoons \text{Zn}^{++} + 2\text{e}^-$ ; the overall reaction is  $\text{Zn} + \text{Cu}^{++} \rightleftharpoons \text{Zn}^{++} + \text{Cu}$ .

The cell voltage is the “ **potential difference**” between the two half cells.

Electrochemists have chosen by convention, the potential of the **standard hydrogen electrode (S.H.E.) as zero**. The redox reaction for this half cell is  $2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{H}_2$ . The **standard potentials,  $E^\circ(\text{S.H.E.})$**  of Cu and Zn are **+0.34** and **-0.76 V** and the cell voltage is,  $E = +0.34 - (-0.76) = +1.10 \text{ V}$ .

(Note: The unit of potential is named volt, in honor of Volta)




Figure 15 - LCD-Clock powered by a tomato battery.



# 1. INTRODUCTION (contd.)

Latest news on S.H.E. : “ ..universal **reference electrode** for which, under standard conditions, the **standard electrode potential** ( $\text{H}^+/\text{H}_2$ ) **is zero** at all temperatures: Last update: **2009-06-16**; <http://goldbook.iupac.org/S05917.html>

There have been attempts in the past to obtain the absolute value of the potential of the S.H.E. (Bard et al [1]). Trasatti (1986) [2] suggested, by theoretical calculations, the value 4.44 (+/-) 0.04V.

Recently, Donald et al (2008) [3] obtained for S.H.E. (by electron capture on gaseous phase nano-drops) the value, 4.20 (+/-) 0.4 V.

The “ **new observation here**” that  $E^\circ$  varies linearly with the ionization potentials (**I**) for many groups of elements, has given the “ **absolute potential of the S.H.E., ( $E^\circ_{I=0}$ )**” (which confirms the value in [3]) (and can be taken as the **final answer**).

Thereby, the “ **absolute redox potentials, have been obtained as the sum,  $E^\circ_{\text{abs}} = E^\circ + E^\circ_{I=0}$** ”, (Heyrovska, [4]).

## 2. LINEAR DEPENDENCE OF AQ. STD. POTENTIALS ( $E^\circ$ ) ON IONIZATION POTENTIALS (I)

The data in [1] have been used to plot  $E^\circ$  vs I in Fig. 1, where I = I (sum) is the sum of the 1st, 2nd, .. ionization potentials, as the case may be, depending on the value of n, ( $\text{ox} + n\text{e}^- = \text{red}$ ).

$$E^\circ = k_{\text{aq}} I - E^\circ_{\text{I}=0}; \text{ absolute aq. redox potential} = E^\circ_{\text{abs}} = E^\circ + E^\circ_{\text{I}=0}$$

$$E^\circ_{\text{I}=0} = \text{Abs. potl. of S.H.E} = 2.87 \text{ V (H/H: halogens) \& 4.20 V (H}^+/\text{H: all others)}$$

(Author: Raji Heyrovská, Brno Conference, 29-30 June 2009)




Fig. 2. Linear dependence of aq. std. potls. on I.

All the straight lines in Fig. 1 follow the simple equation,

$$E^\circ = k_{aq}I - E^\circ_{I=0} ; E^\circ_{abs} = E^\circ + E^\circ_{I=0} (= k_{aq}I) \quad (1a,b)$$

where  $k_{aq}$  is the slope, “  $E^\circ_{I=0}$  is the absolute potential of the S.H.E.” and “  $E^\circ_{abs}$  is the absolute redox potential” .  $E^\circ_{abs} = E^\circ_{I=0}$  when  $E^\circ = 0$  (Eq. 1b).

For Gp. VIIA (halogens,  $(1/2)X_2/X^-$ ),  $E^\circ_{I=0} = 2.87$  (+/-) 0.01 V, and for the others,  $E^\circ_{I=0} = 4.20$  (+/-) 0.03 V, which correspond to the  $(1/2)H_2/H^-$  and  $H^+/H$  redox equilibria resply. Fig. 2 shows plots of  $E^\circ_{abs} (= k_{aq}I)$  vs  $E^\circ$ .

The slopes ( $k_{aq}$ ) of the lines have been used as criteria for classifying the redox couples into groups (i) to (xii) (see: the box in Fig. 1).




Fig. 2 Absolute aq std potentials,  $E^\circ_{abs}$  (calculated as  $k_{aq}I$ ) vs  $E^\circ$ .

### 3. DEPENDENCE OF POLAROGRAPHIC HALF-WAVE POTENTIALS ( $E_{1/2}$ ) ON IONIZATION POTENTIALS (I)

Fig. 3 shows the  $E_{1/2}$  values for **Cu, Pb, Cd, Zn, Mn, Ba** ( $n = 2$  for all) **Al** ( $n = 3$ ) (Heyrovsky, Ilkovic [5a,b]) and **Tl** ( $n = 1$ ) (Heyrovsky, Kuta [6]). The causes for the differences between  $E^\circ$  and  $E_{1/2}$  in some cases (here, **Ba** and **Al**) have been explained by Lingane [7].




Fig 3 Linear dependence of  $E^\circ$  on  $I$ .  
Red filled circles  $E_{1/2}$  values

## ACKNOWLEDGEMENTS

The author thanks the Institute of Biophysics for institutional grant Nos. AV0Z50040507 and AV0Z50040702 of the ASCR and the Organizers of this conference for the opportunity given to present the “ new” scientific results.

## LITERATURE

- Bard, A.J., et al.: Standard potentials in aqueous solutions, Marcel Dekker, 1985, NY.
- Trasatti, S.: Pure & Appl. Chem. 58 (1986) 955.
- Donald, W.A., et al.: J. Am. Chem. Soc. 130 (2008) 3371.
- Heyrovska, R.: submitted
- a) [http://nobelprize.org/nobel\\_prizes/chemistry/laureates/1959/](http://nobelprize.org/nobel_prizes/chemistry/laureates/1959/);
- b) Heyrovsky, J., Ilkovic, D.: Chem. Listy, XXIX (1935) 234; Collection VII (1935) 198.
- 6. Heyrovsky, J., Kuta, J.: Principles of Polarography, Publishing House of the CAS, 1965, Prague.
- 7. Lingane, J.J.: J. Am. Chem. Soc. 61 (1939) 2099.